

**Advisable TOXI FIX ratio per Tonn in mycotoxin contaminated feed**

<b>ANIMAL</b>	<b>% of mould</b>	<b>Ppb presence of toxines*</b>	<b>Toxi fix (kg)</b>
<b>Polutry</b>	<b>Max.60.000</b>	<b>50-100</b>	<b>0,3-1</b>
<b>Pigs</b>	<b>50.000</b>	<b>50-100</b>	<b>0,5-1</b>
<b>Cows, sheeps</b>	<b>80.000</b>	<b>50-100</b>	<b>0,7-1,5</b>
<b>Polutry</b>	<b>Max.120.000</b>	<b>100-200</b>	<b>1-2</b>
<b>Pigs</b>	<b>100.000</b>	<b>100-200</b>	<b>1-2</b>
<b>Cows. Sheeps</b>	<b>150.000</b>	<b>100-200</b>	<b>1,5-2,5</b>
<b>Polutry</b>	<b>Max.240.000</b>	<b>200-750</b>	<b>2-5</b>
<b>Pigs</b>	<b>200.000</b>	<b>200-750</b>	<b>2-6</b>
<b>Cows, sheeps</b>	<b>300.000</b>	<b>200-750</b>	<b>2-7</b>

**\* Aflatoxin, Deoxynivalenol, T-2 toxin, Diacetoscirpenol. In case of Ochratoxin A and Zearalenone half of regular dose is enough.**